

13-gene signature to predict rapid development of brain metastases in HER2-positive advanced breast cancer patients

R Duchnowska¹, J Jassem², C Goswami³, Y Gökmen-Polar³, L Li³,
M Thorat³, A Morimiya³, N Flores⁴, E Hua⁴, S Woditschka⁴, D Palmieri⁴,
SM Steinberg⁴, W Biernat², K Sosińska-Mielcarek⁵, B Szostakiewicz²,
B Czartoryska-Arłukowicz⁶, B Radecka⁷, E Starosławska⁸, Z Tomasevic⁹,
GW Sledge Jr³, PS Steeg⁴, S Badve³

Military Institute of Medicine, Warsaw¹, Poland, Medical University,
Gdansk, Poland², Indiana University School of Medicine, Indianapolis, IA
USA³, National Cancer Institute, Bethesda, MD USA⁴, Regional Oncology
Center, Gdansk, Poland⁵, Oncology Center, Białystok, Poland⁶, Oncology
Center, Opole, Poland⁷, Oncology Center, Lublin, Poland⁸, Institute for
Oncology
and Radiology, Belgrade, Serbia⁹

**Data was partially presented at the
ASCO Annual Meeting in 2008 and 2012**

Background and Study Aims

- Brain metastases of breast cancer constitute an important part of therapeutic failures and are associated with severe morbidity and mortality
- The risk of brain metastases is particularly high in HER2-positive advanced breast cancer patients
- Using RNA-microarray technology, we earlier developed in this group a 13-gene signature strongly predicting for rapid development of brain metastases (J Clin Oncol 2008; 26: 45s)
- Now, using qRT-PCR technology we investigated the clinical outcome of genes constituting this signature in an independent group of patients, and employed a culture model system

Discovery (DASL) Series: HER2-Positive Advanced Breast Cancer Patients (n=87)

Age	Mean, years (range)	49 (29-75)
Menopausal status	Pre/Post/Unknown	47/39/1
Pathology type	Ductal/Lobular/Other	69/4/14
Dominant site of disease	Viscera/Soft tissue/ Bones	78/4/5
ER status	(+)/(+)/Unknown	29/55/3
Grade	2/3/Unknown	26/48/13
Trastuzumab therapy	Yes/No	72/15
Brain metastases (BM)	Yes/No	47/40
Median time to BM development	55 months	

DASL Analysis

- RNA (200 ng) extracted from FFPE primary tumor samples using HighPure RNA Paraffin Kit (Roche Applied Bioscience)
- RNA pre-qualified using iScript (Bio-Rad) to reverse transcribe and SYBR Green Master Mix (Applied Biosystems) to perform quantitative RT-PCR
- DASL assay of 502 known cancer genes performed using the Sentrix Universal Array (Illumina Corp)
- Microarray data normalized through the sample median
- Biostatistics data analysis using t-test with unequal variances, significant analysis of microarray, and predictive analysis of microarray

13-gene Signature

HER2 related

- *CDK4* (Cyclin dependent kinase 4)
- *CCNC* (Cyclin C)
- *PTK2* (FAK, Focal adhesion kinase)
- *MYC* (Myc)


DNA double strand break repair

- *BARD1*
- *RAD 51* (Rad 51)
- *FANCG* (Fanconia anemia group G)


Others

- *PCNA* (Proliferating cell nuclear antigen)
- *PRCC* (Papillary renal cell carcinoma-translocation associated)
- *TPR* (Translocated promoter region)
- *EMS1* (Cortactin)
- *DSP* (Desmoplakin)
- *HDGF* (Hepatocyte derived growth factor)

Time from Diagnosis to Brain Relapse by 13-Gene Signature (DASL Series)



Time from Diagnosis to Brain Relapse by 13-Gene Signature in ER-, HER2+ primary tumors (DASL Series)



Independent qRT-PCR Series: HER2-Positive Advanced Breast Cancer Patients (n=75)


Age	Mean, years (range)	52 (28-71)
Menopausal status	Pre/Post/Unknown	47/28/0
Pathology type	Ductal/Lobular/Other+Unkn.	58/5/12
Dominant site of disease	Viscera/Soft tissue/Bones	52/15/7
ER status	(+)/(-)/Unknown	32/42/1
Grade	2/3/Unknown	32/33/10
Trastuzumab therapy	Yes/No	63/12
Brain metastases (BM)	Yes/No	34/41
Median time to BM development	65 months	

qRT-PCR Cohort Analysis


Statistical Methods

- qRT-PCR was performed for 13 genes plus two control genes (*GAPDH*, *ACTB*)
- Delta cycle threshold (ΔCt) values were generated by ABI Biosystems Data Assist Software and normalized to control genes
- ΔCt values were standardized for use in univariate Cox regressions by subtraction of mean, division by SD for each gene
- ΔCt values were weighted by individual Cox PH coefficients
- Expression score = sum of weighted ΔCt values for the 13 genes
- Median of data used to divide into high vs. low expression groups
- Cox regression used to test association of expression score with time to brain metastasis

Time from Diagnosis to Brain Relapse by 13-Gene Signature (qRT-PCR Series)



Time from Diagnosis to Brain Relapse by 13-Gene Signature in ER-, HER2+ Primary Tumors (qRT-PCR Series)



Association Between 13-Gene Signature and Standard Patient Traits

	Number (%) of patients with 13-gene signature					
	DASL cohort			PCR cohort		
	Low	High	P	Low	High	P
ER Negative	24 (28%)	30 (35%)		18 (24%)	24 (32%)	
Positive	18 (21%)	13 (15%)	0.26	19 (26%)	13 (18%)	0.24
PR Negative	31 (36%)	30 (35%)		25 (34%)	28 (38%)	
Positive	11 (13%)	13 (15%)	0.81	12 (16%)	9 (12%)	0.61
Menopausal status Pre	25 (29%)	20 (23%)		15 (20%)	13 (17%)	
Post	17 (20%)	24 (28%)	0.20	23 (31%)	24 (32%)	0.81
Primary tumor grade 2	20 (25%)	15 (19%)		19 (29%)	13 (20%)	
3	18 (23%)	26 (33%)	0.18	12 (19%)	21 (32%)	0.08
Patient age (median (years))	50	48.5	0.94	53	51	0.55

DNA Double Strand Break Repair Genes in the 13-Gene Signature (Repair by Homologous Recombination)




- *RAD51*; serves to recombine the broken single stranded DNA with the homologous, intact strand
- *BARD1*; an obligate binding partner for *BRCA1*
- *FANCG*; a part of the Fanconi Anemia complex


MCF-10A Three Dimensional Culture Model

- Does overexpression of *RAD51* and/or *BARD1* exert a phenotypic effect on breast tumorigenesis?

MCF-10A three dimensional culture model



Normal breast acinus




HER2 overexpression


Nature Rev. Cancer 5:675,2005

- Does overexpression of *RAD51* or *BARD1* promote brain metastasis?
Experimental brain metastasis assays ongoing


Transfection of MCF-10A Immortal Breast Cells With *RAD51*, *BARD1* and/or HER2



In three-dimensional culture, the transfectants developed distinct morphologies:



***RAD51* Overexpression in Immortal MCF-10A Cells Increased Invasion In Vitro**



Fold change (FC) of % invasive acini growth for each transfected gene in MCF10A compared with MCF10A control transfectants

Conclusions

- 13-gene signature and ER-negativity predict rapid development of brain metastases in HER2-positive advanced breast cancer patients
- *RAD51* may promote aggressiveness in breast epithelial cells
- These data may be useful in the design of brain metastasis preventive trials and may prompt new treatment strategies

Acknowledgments

Special thanks to all pathologists
who provided archive tumor samples
for molecular analysis